

On the Geometry of Bäcklund Transformations

M. Palese and E. Winterroth*

Department of Mathematics, University of Torino

Via C. Alberto 10, 10123 Torino, Italy

e-mails: PALESE@DM.UNITO.IT, EKKEHART@DM.UNITO.IT

Abstract

The geometry of an admissible Bäcklund transformation for an exterior differential system is described by an admissible Cartan connection for a geometric structure on a tower with infinite-dimensional skeleton which is the universal prolongation of a $|1|$ -graded semi-simple Lie algebra.

2000 MSC: 14M17, 53C05, 53C10, 53C30, 58J70, 58J72, 58A15.

Key words: generalized homogeneous spaces, Bäcklund transformations, connections.

1 Towers on skeletons

In the following we consider infinite-dimensional objects in the $\mathfrak{p}$ -category of objects obtained as *projective limits* of finite-dimensional ones [3].

Definition 1 An algebraic skeleton on a finite-dimensional vector space V is a triple (E, G, ρ) , with G a $\mathfrak{p}$ -Lie group, $E = V \oplus \mathfrak{g}$, $\mathfrak{g}$ the Lie algebra of G , and ρ a representation of G on E such that $\rho(g)x = Ad(g)x$, for $g \in G$, $x \in \mathfrak{g}$. An infinitesimal skeleton can be analogously defined *via* the representation of $\mathfrak{g}$ on E . $\square$

Definition 2 Let (E, G, ρ) be a skeleton on V and Z a manifold of type V [3]. We say that a $\mathfrak{p}$ -principal fibre bundle $P(Z, G)$ provided with an absolute parallelism ω on P is a *tower* on Z with skeleton (E, G, ρ) if ω takes values in E and satisfies: $R_g^*\omega = \rho(g)^{-1}\omega$, for $g \in G$; $\omega(\tilde{A}) = A$, for $A \in \mathfrak{g}$; here R_g denotes the right translation and $\tilde{A}$ the fundamental vector field induced on P from A . $\square$

*Both of them supported by GNFM of INdAM and University of Torino proj. *Giovani Ricercatori 2001*. E.W. partially supported also by University of Erlangen–Nürnberg and CNR grant n. 201.21-00.01.01.

1.1 Cartan connections

Let $\mathfrak{g}$ be a Lie algebra and $\mathfrak{k}$ a Lie subalgebra of $\mathfrak{g}$. Let K be a Lie group with Lie algebra $\mathfrak{k}$ equipped with a representation $Ad : K \rightarrow GL(\mathfrak{g})$ such that its differential coincides with the adjoint representation of $\mathfrak{k}$ on $\mathfrak{g}$.

Definition 3 Let $P(Z, K)$ be a principal fibre bundle over a manifold Z with structure group K . A *Cartan connection* in P of type $(\mathfrak{g}, K)$ is a 1-form ω on P with values in $\mathfrak{g}$ satisfying the following conditions: $\omega|_{T_u P} : T_u P \rightarrow \mathfrak{g}$ is an isomorphism $\forall u \in P$; $R_g^* \omega = Ad(g)^{-1} \omega$ for $g \in K$; $\omega(\tilde{A}) = A$ for $A \in \mathfrak{k}$. $\square$

$(\mathfrak{g}, K, Ad)$ is a skeleton on V , with $\mathfrak{g} = \mathfrak{k} \oplus V$. Then it is clear that a Cartan connection (P, Z, K, ω) of type $(\mathfrak{g}, K)$ is a tower on Z .

In the following we assume the Lie algebra $\mathfrak{g}$ to be a generalized semi-simple $|1|$ -graded Lie algebra *i.e.* $\mathfrak{g} = \mathfrak{g}_{-1} \oplus \mathfrak{g}_0 \oplus \mathfrak{g}_1$ [4].

Remark 1 According with the Lie algebra $|1|$ -grading the Cartan connection form ω and its curvature κ split as $\omega = \omega_{-1} \oplus \omega_0 \oplus \omega_1$ and $\kappa = \kappa_{-1} \oplus \kappa_0 \oplus \kappa_1$. $\square$

Definition 4 Let G be a semi-simple Lie group, with $|1|$ -graded Lie algebra $\mathfrak{g}$ as above and K the closed subgroup of G corresponding to the Lie algebra $\mathfrak{g}_0 \oplus \mathfrak{g}_1$. A K -structure on Z is a principal fiber bundle $P \rightarrow Z$ with structure group K equipped with a *soldering* form $\theta = \theta_{-1} \oplus \theta_0 \in \Omega^1(P, \mathfrak{g}_{-1} \oplus \mathfrak{g}_0)$ such that: $\theta_{-1}(\xi) = 0$, if and only if ξ is a vertical vector; $\theta_0(\tilde{X} + \tilde{Z}) = Y$, $\forall Y \in \mathfrak{g}_0, Z \in \mathfrak{g}_1$; $(R_b)^* \theta = Ad(b^{-1})\theta$, $\forall b \in K$, where Ad means the action on the vector space $\mathfrak{g}_{-1} \oplus \mathfrak{g}_0 \simeq \mathfrak{g}/\mathfrak{g}_1$ induced by the adjoint action. $\square$

Definition 5 Let (P, θ) be a K -structure on Z . A Cartan connection ω on P is called *admissible* if and only if it is of the form $\omega = \theta_{-1} \oplus \theta_0 \oplus \omega_1$. $\square$

Remark 2 If the K -structure has zero torsion, *i.e.* if it is a reduction of $L^2(Z) \rightarrow Z$ to K , then the curvature of the induced Cartan connection $\omega = \theta_{-1} \oplus \theta_0 \oplus \omega_1$ is such that $\kappa_{-1} = 0$ [2]. $\square$

2 Bäcklund transformations and induced Cartan connections

Let $\pi : U \rightarrow X$, $\tau : Z \rightarrow X$, be two (vector) bundles with local fibered coordinates (x^α, u^A) and (x^α, z^i) , respectively, where $\alpha = 1, \dots, m = \dim X$, $A = 1, \dots, n = \dim U - \dim X$, $i = 1, \dots, N = \dim Z - \dim X$. A system of non-linear field equations of order k on U is geometrically described as an exterior differential system ν on $J^k U$. The solutions of the field equations are (local) sections σ of $U \rightarrow X$ such that $(j^k \sigma)^* \nu = 0$. We shall also denote by $J^\infty \nu$ (*resp.* $j^\infty \sigma$) the infinite order jet prolongation of ν (*resp.* σ).

2.1 Admissible Bäcklund transformations

Let B be the infinite-order contact transformations group on $J^\infty U$.

Definition 6 The group of (infinitesimal) *Bäcklund transformations for the system ν* is the closed subgroup $\tilde{K}$ of B which leaves invariant solution submanifolds of $J^\infty \nu$. The group of (infinitesimal) *generalized Bäcklund transformations for the system ν* is the closed subgroup K of B which leaves invariant $J^\infty \nu$ [1]. $\square$

Let $\pi : U \rightarrow X$, $\tau : Z \rightarrow X$, be vector bundles as the above and $\pi^1 : J^1 U \rightarrow X$, $\tau^1 : J^1 Z \rightarrow X$, the first order jet prolongations bundles, with local fibered coordinates $(x^\alpha, u^A, u_\alpha^A)$, $(x^\alpha, z^i, z_\alpha^i)$, respectively. Furthermore, let $(\partial_\beta, \partial_A, \partial_A^\beta)$, $(\partial_\beta, \partial_i, \partial_i^\beta)$ and $(dx^\beta, du^A, du_\beta^A)$, $(dx^\beta, dz^i, dz_\beta^i)$ be local bases of tangent vector fields and 1-forms on $J^1 U$ and $J^1 Z$, respectively.

Definition 7 We define a Bäcklund map to be the fibered morphism over Z : $\phi : J^1 U \times_X Z \rightarrow J^1 Z : (x^\alpha, u^A, u_\alpha^A; z^i) \mapsto (x^\alpha, z^i, z_\alpha^i)$, with $z_\alpha^i = \phi_\alpha^i(x^\beta, u^A, u_\beta^A; z^j)$.

The fibered morphism ϕ is said to be an *admissible* Bäcklund transformation for the differential system ν if $\phi_\alpha^i = \mathcal{D}_\alpha \phi^i$ and the integrability conditions coincide with the exterior differential system ν . $\square$

Remark 3 By pull-back of the contact structure on $J^1 Z$, the Bäcklund morphism induces an horizontal distribution, the *induced Bäcklund connection*, on the bundle $(J^1 U \times_X Z, J^1 U, \pi_0^*(\eta))$ [6]. $\square$

Theorem 1 *The following statements are equivalent [5].*

1. ϕ is an admissible Bäcklund transformation for the differential system ν .
2. The induced Bäcklund connection is $\tilde{K}$ -invariant, where $\tilde{K}$ is a normal subgroup $\tilde{K} \subset (\tilde{K} \cap K) \subset B$ leaving invariant (the infinite order prolongation of) ν and its solutions.

Let now Z be a vector bundle over the basis X , the fibers of which are modelled over the homogeneous space $\tilde{K}/\tilde{K}$ such that fibers are vector spaces of type $\tilde{\mathfrak{K}}_- = \oplus_{p < 0} \tilde{\mathfrak{K}}_p$, with $\tilde{\mathfrak{K}}_-$ a graded abelian Lie algebra. Let P be a tower on Z with algebraic skeleton $(\tilde{\mathfrak{k}}, \tilde{\mathfrak{K}}, Ad)$, where $\tilde{\mathfrak{k}}$ is the Lie algebra of $\tilde{K}$. Suppose U be a vector bundle (over the same basis X) with a left action λ of $\tilde{K}$ on U (as a manifold). For each tower $(P, Z, \tilde{K}, \omega)$ we have a vector bundle $U_\lambda(Z) = P \times_{\tilde{K}} U$ over X and *vice versa*.

Assume $\tilde{\mathfrak{k}}$ to be the universal prolongation [7] of a $|1|$ -graded semi-simple Lie algebra $\mathfrak{g}$ such that $\tilde{\mathfrak{k}}$ and $\tilde{\mathfrak{K}}_-$ are (the prolongation of) $\mathfrak{g}_0 \oplus \mathfrak{g}_1$ and $\mathfrak{g}_{-1}$, respectively.

Theorem 2 *A Bäcklund transformation admissible for an exterior differential system induces the tower $(P, Z, \tilde{K}, \omega)$, where ω is an admissible Cartan connection for a $\tilde{K}$ -structure over Z .*

PROOF. It follows from Remark 2 and Theorem 1. In fact, a Bäcklund morphism can be seen as a $\bar{K}$ -equivariant section of the bundle $U_\lambda(Z) \rightarrow Z$; it induces a reduction of $L^2(Z) \rightarrow Z$ defining a $\bar{K}$ -structure on Z with zero torsion. The admissible Cartan connection is the tower $(P, \bar{K}, \theta)$ induced [2] from a $\mathfrak{g}_0$ -principal connection on the underlying first order structure. QED

As a consequence one can built a cohomological theory of complete integrability for (nonlinear) exterior differential systems. Cohomological conditions are in fact given for a graded simple Lie algebra to be a universal prolongation [7]. This topic will be developed elsewhere.

Acknowledgments

Thanks are due to M. Francaviglia, R.A. Leo and G. Soliani for useful discussions.

References

- [1] R.L. Anderson and N.H. Ibragimov, *Lie-Bäcklund Transformations in Applications*, SIAM, Philadelphia 1979
- [2] A. Čap, J. Slovák and V. Souček, *Acta Math. Univ. Comenian. (N.S.)* **66**, 33 (1997).
- [3] T. Morimoto, *Hokkaido Math. Jour.* **22**, 263 (1993).
- [4] T. Ochiai, *Trans. Amer. Math. Soc.* **152**, 159 (1970).
- [5] M. Palese and E. Winterroth, *Phys. Lett. B* **532**, 129 (2002).
- [6] F.A.E. Pirani, D.C. Robinson, W.F. Shadwick, *Local Jet Bundle Formulation of Bäcklund Transformations*, Math. Phys. Stud., D. Reidel Publishing Company, Dordrecht, Holland, 1979.
- [7] N. Tanaka, *Hokkaido Math. Jour.* **8**, 23 (1979); *J. Math. Soc. Japan* **19**, 215 (1967).